

Connecting you to the support you deserve

Increased risk of death in elderly patients with dementia-related psychosis: Medicines like NUPLAZID® (pimavanserin) can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). NUPLAZID is not approved for the treatment of patients with dementia-related psychosis unrelated to the hallucinations and delusions associated with Parkinson's disease psychosis.

Indication

NUPLAZID is a prescription medicine used to treat hallucinations and delusions associated with Parkinson's disease psychosis.

See additional Important Safety Information including **Boxed WARNING** on back cover. Please read the accompanying full Prescribing Information.

A clear path forward

Beginning a new medication can feel overwhelming and confusing for you and your loved ones. NUPLAZIDconnect™ was specifically created to help you get the support you deserve for a clear path forward.

Your dedicated NUPLAZIDconnect Care Coordinator will connect you with the answers you need so you have a clear understanding of what to expect.

“[NUPLAZIDconnect] answered questions that I didn’t even know to ask. I cried—this was exactly what I needed.

CAREGIVER & DAUGHTER OF A LOVED ONE
LIVING WITH HALLUCINATIONS AND DELUSIONS
ASSOCIATED WITH PARKINSON’S DISEASE



Your Care Coordinator can help you in a variety of ways, including:



Navigating insurance and managing costs

Your Care Coordinator can help explain your insurance coverage and the financial support options available to you.



Getting your prescription filled

Your Care Coordinator will work with your specialty pharmacy to fill your NUPLAZID (primavanserin) prescription. They’ll make sure you understand when and how your medication will be delivered to you.



Answering your questions

Your Care Coordinator can also connect you to a Nurse Educator to answer questions you may have about your medication, such as:

- When and how to take NUPLAZID
- What side effects you may experience
- How long it may take for NUPLAZID to start working

Please don’t hesitate to reach out—we’re here for you and look forward to helping you find a clear path forward with NUPLAZIDconnect



**Call 1-844-737-2223, Monday through Friday 8:30 AM–8:30 PM EST,
or visit [NUPLAZIDconnect.com](https://www.nuplazidconnect.com)**

See Important Safety Information including **Boxed WARNING** on back cover. Please read the accompanying full Prescribing Information.

Indication

NUPLAZID is a prescription medicine used to treat hallucinations and delusions associated with Parkinson's disease psychosis.

Important Safety Information for NUPLAZID (pimavanserin)

Increased risk of death in elderly patients with dementia-related psychosis: Medicines like NUPLAZID can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). NUPLAZID is not approved for the treatment of patients with dementia-related psychosis unrelated to the hallucinations and delusions associated with Parkinson's disease psychosis.

Do not take NUPLAZID if you have had an allergic reaction to any of the ingredients in NUPLAZID. Allergic reactions have included rash, hives, swelling of the tongue, mouth, lips, or face, throat tightness, and shortness of breath.

NUPLAZID may cause serious side effects including:

QT Interval Prolongation: NUPLAZID may increase the risk of changes to your heart rhythm. This risk may increase if NUPLAZID is taken with certain other medications known to prolong the QT interval. **Tell your healthcare provider about all the medicines you take or have recently taken.**

Do not take NUPLAZID if you have certain heart conditions that change your heart rhythm. It is important to talk to your healthcare provider about this possible side effect. Call your healthcare provider if you feel a change in your heartbeat.

Tell your healthcare provider about all the medicines you take. Other medicines may affect how NUPLAZID works. Some medicines should not be taken with NUPLAZID. Your healthcare provider can tell you if it is safe to take NUPLAZID with your other medicines. Do not start or stop any medicines while taking NUPLAZID without talking to your healthcare provider first.

The **most common side effects** of NUPLAZID include swelling in the legs or arms, nausea, confusion, hallucination, constipation, and changes to normal walking. These are not all the possible side effects of NUPLAZID. For more information, ask your healthcare provider about this medicine.

It is not known if NUPLAZID is safe and effective in people under 18 years of age.

Dosage and Administration

The recommended dose of NUPLAZID is 34 mg once per day, taken by mouth.

Please read the accompanying full Prescribing Information.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You can also call ACADIA Pharmaceuticals Inc. at 1-844-4ACADIA (1-844-422-2342).

NUPLAZID is available as 34 mg capsules, 17 mg tablets, and 10 mg tablets.

NUPLAZID® and NUPLAZIDconnect™ are registered trademarks of ACADIA Pharmaceuticals Inc. ©2018 ACADIA Pharmaceuticals Inc. All rights reserved. NC-0131 07/18.