


Important Information for
Patients and Caregivers

ONCE-DAILY
NUPLAZID[®]
(pimavanserin) 34mg capsules

GETTING STARTED WITH NUPLAZID



Please read this important information before using NUPLAZID[®] (pimavanserin) and each time you get a refill because this information may change. This information does not take the place of talking to your doctor about your medical condition or your treatment.

Important Safety Information for NUPLAZID (pimavanserin)

Increased risk of death in elderly patients with dementia-related psychosis: Medicines like NUPLAZID can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). NUPLAZID is not approved for the treatment of patients with dementia-related psychosis unrelated to the hallucinations and delusions associated with Parkinson's disease psychosis.

ABOUT NUPLAZID



What is NUPLAZID® (pimavanserin) used for?

Some people may begin to see or hear things that others don't (hallucinations) and/or may believe things that are not true (delusions). This may occur over the course of their Parkinson's disease. NUPLAZID (new-PLA-zid) is specifically used to treat these non-motor symptoms in patients with Parkinson's disease.



How do I take NUPLAZID?

It is recommended that you take one 34 mg capsule of NUPLAZID once daily by mouth. Some medicines may interact with NUPLAZID so your healthcare provider may prescribe a different dose of NUPLAZID.



How long does it take for NUPLAZID to work?

NUPLAZID may begin to reduce hallucinations and delusions as soon as 4 weeks, but it may take up to 6 weeks to feel the full effect. You should speak with your healthcare provider before you stop taking NUPLAZID.



Can I take NUPLAZID with other Parkinson's disease medicines?

You can take NUPLAZID without adjusting doses of carbidopa/levodopa medicines. Some medicines should not be taken with NUPLAZID. **Tell your healthcare provider about all the medicines you are taking.**

Additional Important Safety Information



Who should not take NUPLAZID?

Do not take NUPLAZID if you have had an allergic reaction to any of the ingredients in NUPLAZID. Allergic reactions have included rash, hives, swelling of the tongue, mouth, lips, or face, throat tightness, and shortness of breath.



What additional warnings should I know about NUPLAZID?

NUPLAZID may increase the risk of changes to your heart rhythm. This risk may increase if NUPLAZID is taken with certain other medications known to prolong the QT interval.

Tell your healthcare provider about all the medicines you take or have recently taken.

Do not take NUPLAZID if you have certain heart conditions that change your heart rhythm. It is important to talk to your healthcare provider about this possible side effect. Call your healthcare provider if you feel a change in your heartbeat.





What other medicines might interact with NUPLAZID?

Tell your healthcare provider about all the medicines you take. Other medicines may affect how NUPLAZID works. Some medicines should not be taken with NUPLAZID. Your healthcare provider can tell you if it is safe to take NUPLAZID with your other medicines. Do not start or stop any medicines while taking NUPLAZID without talking to your healthcare provider first.



What are the common side effects of NUPLAZID?

The **most common side effects** of NUPLAZID include swelling in the legs or arms, nausea, confusion, hallucination, constipation, and changes to normal walking. These are not all the possible side effects of NUPLAZID. For more information, ask your healthcare provider about this medicine.





How do I get NUPLAZID?

Start with NUPLAZIDconnect[™]. Beginning a new medication can feel overwhelming and confusing for you and your loved ones. NUPLAZIDconnect was specifically created to help you get the support you deserve for a clear path forward.

Your dedicated NUPLAZIDconnect Care Coordinator will connect you with the answers you need so you have a clear understanding of what to expect.

Your Care Coordinator can help you in a variety of ways, including:

Navigating insurance and managing costs

- Your Care Coordinator can help explain your insurance coverage and the financial support options available to you

Getting your prescription filled

- Your Care Coordinator will work with your specialty pharmacy to fill your NUPLAZID prescription. They'll make sure you understand when and how your medication will be delivered to you

Answering your questions

- Your Care Coordinator can also connect you to a Nurse Educator to answer questions you may have about your medication, such as:
 - When and how to take NUPLAZID
 - What side effects you may experience
 - How long it may take for NUPLAZID to start working

ONCE-DAILY
NUPLAZID[®]
(pimavanserin) 34mg capsules

Questions?

Call **1-844-737-2223**,
Monday through Friday,
8:30 AM - 8:30 PM EST.

NUPLAZIDconnect.com

NUPLAZID
connect[™]

Additional Important Safety Information

It is not known if NUPLAZID is safe and effective in people under 18 years of age.

This is not the complete safety information. For full safety information, please read the attached full [Prescribing Information](#), including **Boxed WARNING**, or visit NUPLAZID.com.

If you have any questions about this medicine or this information, be sure to discuss them with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You can also call ACADIA Pharmaceuticals Inc. at 1-844-4ACADIA (1-844-422-2342).